

Synthesis and cytostatic evaluation of some 2-(5-substituted-2-oxoindolin-3-ylidene)-N-substituted hydrazine carbothioamide

Subhas S. Karki, Amol Kulkarni, Nishith Teraiya, Erik De Clercq & Jan Balzarini

Medicinal Chemistry Research

ISSN 1054-2523

Volume 20

Number 8

Med Chem Res (2011) 20:1229-1234

DOI 10.1007/s00044-010-9458-3



Your article is protected by copyright and all rights are held exclusively by Springer Science+Business Media, LLC. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your work, please use the accepted author's version for posting to your own website or your institution's repository. You may further deposit the accepted author's version on a funder's repository at a funder's request, provided it is not made publicly available until 12 months after publication.

Synthesis and cytostatic evaluation of some 2-(5-substituted-2-oxoindolin-3-ylidene)-*N*-substituted hydrazine carbothioamide

Subhas S. Karki · Amol Kulkarni · Nishith Teraiya ·
Erik De Clercq · Jan Balzarini

Received: 16 May 2010 / Accepted: 25 September 2010 / Published online: 19 October 2010
© Springer Science+Business Media, LLC 2010

Abstract Various substituted 2-(5-substituted-2-oxoindolin-3-ylidene)-*N*-substituted hydrazine carbothioamide **4a–g** and 2-(5-substituted-1-(4-substituted benzyl)-2-oxoindolin-3-ylidene)-*N*-substituted hydrazine carbothioamide **5a–k** were synthesized. The compounds were evaluated for their cytostatic activity against human Molt4/C8 and CEM T-lymphocytes as well as murine L1210 leukemia cells. Several of these compounds were endowed with low micromolar 50%-inhibitory concentration (IC₅₀) values, and some were virtually equally potent as melphalan. The most potent inhibitors against the murine leukemia cells were also most inhibitory against human T-lymphocyte tumor cells. 2-(5-fluoro-1-(4-fluorobenzyl)-2-oxoindolin-3-ylidene)-*N*-*p*-tolylhydrazine carbothioamide (**5b**) emerged as the most potent cytostatic compound among the tested compounds. The encouraging cytostatic data provide an adequate rationale for further modification of these molecular scaffolds.

Keywords 2,3-dioxo-2,3-dihydroindole ·
Thiosemicarbazones · Cytostatic assays

Introduction

Isatin has been known for about 150 years and has recently been found, like 2,3-dioxo-indoles and endogenous poly-functional heterocyclic compounds, to exhibit biological activity in mammals (Somogyi, 2001). Isatin is also a synthetically versatile substrate that can be used to prepare a large variety of heterocyclic compounds, such as indoles and quinolines, and as a raw material for drug synthesis (Silva *et al.*, 2001). Schiff bases and Mannich bases of isatin are known to possess a wide range of pharmacological properties including antibacterial (Pandeya *et al.*, 1998; Sarangapani and Reddy, 1994; Varma and Nobles, 1975), anticonvulsant (Sridhar *et al.*, 2002; Varma *et al.*, 2004), anti-HIV (Pandeya *et al.*, 1998 1999a, b 2000; Sriram *et al.*, 2000), antifungal (Pandeya *et al.*, 1999a, b), antiviral (Singh *et al.*, 1983), and anticancer activity (Karki *et al.*, 2004, 2007, 2009).

Thiosemicarbazones of various aldehydes and ketones occupy a special place among organic ligands, since they contain various donor atoms and are able to change density depending on the starting reagents and their reaction conditions. Isatin-3-thiosemicarbazones (1H-indole-2,3-dioxo-3-thiosemicarbazones) have been studied extensively due to their important biological activities (Karki *et al.*, 2009), since 1-methylisatin-3-thiosemicarbazone (Marboran) was found to be active in the treatment of smallpox. Previous studies by our group have revealed the promising cytotoxic properties of various 2,3-dioxo-2,3-dihydroindole thiosemicarbazones (Karki *et al.*, 2009). Therefore, we have performed the synthesis of new *N*-4-aryl thiosemicarbazone derivatives of substituted 2,3-dioxo-2,3-dihydroindoles and studied their antiproliferative activity against human Molt 4/C8 and CEM T-lymphocytes as well as murine L1210 leukemia cells.

S. S. Karki (✉) · A. Kulkarni · N. Teraiya
Department of Pharmaceutical Chemistry, KLE University's
College of Pharmacy, Rajajinagar, Bangalore 560010,
Karnataka, India
e-mail: subhasskarki@gmail.com

E. De Clercq · J. Balzarini
Rega Institute for Medical Research, Katholieke Universiteit
Leuven, Minderbroedersstraat 10, B-3000 Leuven, Belgium

Experimental

Chemistry

All reagents were obtained from Sigma-Aldrich Mumbai, and Loba Chemie, Mumbai. All the solvents used in these studies were dried and distilled before use. Melting points were recorded on a Veego VMP-PM digital melting point apparatus, and are uncorrected. FT-IR spectras were recorded on Shimadzu 8400 S, FT-IR. ^1H NMR spectra were recorded on 300 MHz JEOL NMR Spectrophotometer in CDCl_3 and $\text{DMSO}-d_6$. All spectras were obtained from Pune University, Maharashtra, India.

Synthesis of the intermediate indoline-2,3-diones

The synthesis of the intermediate indoline-2,3-diones was accomplished using a literature methodology (Marvel and Heirs, 1941), and a previously reported procedure was used to convert these compounds to the corresponding 1-benzylindoline-2,3-diones (Azizian *et al.*, 2003). The *N*-4-arylthiosemicarbazides required in the preparation of **4a–g** and **5a–k** were prepared by a literature methodology (Sen and Sengupta, 1962; Lieber *et al.*, 1957).

General procedure for the synthesis of **4a–g** and **5a–k**

A mixture of the indoline-2,3-diones (**2**)/1-benzylindoline-2,3-dihydroindole (**3**) (0.005 mol), *N*-(4-substituted aryl)-thiosemicarbazides (0.005 mol), acetic acid (0.5–1.0 ml), and ethanol (100 ml) was heated under reflux until the reaction was completed (approximately 4 h). Approximately half of the ethanol was removed in vacuo, and the solution was left overnight at room temperature. The solid which precipitated was collected by filtration, washed with cold ethanol, and recrystallized from ethanol:chloroform (9:1) to give compounds **4a–g** and **5a–k**, respectively.

2-(5-chloro-2-oxoindolin-3-ylidene)-*N*-(4-chlorophenyl)hydrazine carbothioamide (**4a**)

Yield 69%. mp 260–261°C. FT-IR ν_{max} cm^{-1} (KBr): 1129, 1289, 1686, 3061, 3143, 3226, 3246. ^1H -NMR δ : 6.9–7.8 (m, 7H, Ar-H), 10.94 (s, 1H, CO-NH), 11.38 (s, 1H, N-NH), 12.66 (s, 1H, NH).

2-(5-bromo-2-oxoindolin-3-ylidene)-*N*-(4-chlorophenyl)hydrazine carbothioamide (**4b**)

Yield 86%. mp 255–258°C. FT-IR ν_{max} cm^{-1} (KBr): 1131, 1295, 1672, 3019, 3113, 3220, 3250. ^1H -NMR δ : 6.9–8.0 (m, 7H, Ar-H), 10.94 (s, 1H, CO-NH), 11.38 (s, 1H, N-NH), 12.66 (s, 1H, NH).

2-(5-bromo-2-oxoindolin-3-ylidene)-*N*-*p*-tolylhydrazine carbothioamide (**4c**)

Yield 76%. mp 255–260°C. FT-IR ν_{max} cm^{-1} (KBr): 1184, 1315, 1692, 3015, 3123, 3212, 3248. ^1H -NMR δ : 2.51 (s, 3H, CH_3), 6.9–8.0 (m, 7H, Ar-H), 10.86 (s, 1H, CO-NH), 11.36 (s, 1H, N-NH), 12.59 (s, 1H, NH).

2-(5-fluoro-2-oxoindolin-3-ylidene)-*N*-*p*-tolylhydrazine carbothioamide (**4d**)

Yield 78%. mp 276–280°C. FT-IR ν_{max} cm^{-1} (KBr): 1154, 1291, 1682, 2995, 3116, 3234, 3256. ^1H -NMR δ : 2.51 (s, 3H, CH_3), 6.9–7.6 (m, 7H, Ar-H), 10.81 (s, 1H, CO-NH), 11.27 (s, 1H, N-NH), 12.65 (s, 1H, NH).

2-(6-chloro-5-fluoro-2-oxoindolin-3-ylidene)-*N*-(4-chlorophenyl)hydrazine carbothioamide (**4e**)

Yield 70%. mp 278–280°C. FT-IR ν_{max} cm^{-1} (KBr): 1164, 1317, 1698, 3019, 3133, 3213, 3243. ^1H -NMR δ : 6.9–7.8 (m, 6H, Ar-H), 10.2 (s, 1H, CO-NH), 11.54 (s, 1H, N-NH), 12.63 (s, 1H, NH).

2-(6-chloro-5-fluoro-2-oxoindolin-3-ylidene)-*N*-*p*-tolylhydrazine carbothioamide (**4f**)

Yield 69%. mp 267–270°C. FT-IR ν_{max} cm^{-1} (KBr): 1180, 1310, 1672, 3010, 3128, 3215, 3237. ^1H -NMR δ : 3.27 (s, 3H, CH_3), 6.9–7.8 (m, 6H, Ar-H), 10.14 (s, 1H, CO-NH), 11.52 (s, 1H, N-NH), 12.62 (s, 1H, NH).

N-(4-chlorophenyl)-2-(5-fluoro-2-oxoindolin-3-ylidene)hydrazine carbothioamide (**4g**)

Yield 73%. mp 290–293°C. FT-IR ν_{max} cm^{-1} (KBr): 1165, 1295, 1682, 3008, 3121, 3210, 3238. ^1H -NMR δ : 6.6–7.8 (m, 7H, Ar-H), 11.10 (s, 1H, CO-NH), 11.29 (s, 1H, N-NH), 12.52 (s, 1H, NH).

N-(4-chlorophenyl)-2-(5-fluoro-1-(4-fluorobenzyl)-2-oxoindolin-3-ylidene)hydrazine carbothioamide (**5a**)

Yield 74%. mp 197–200°C. FT-IR ν_{max} cm^{-1} (KBr): 1119, 1298, 1696, 3042, 3129, 3214, 3257. ^1H -NMR δ : 5.0 (s, 2H, CH_2), 7.0–7.7 (m, 11H, Ar-H), 10.96 (s, 1H, N-NH), 12.65 (s, 1H, NH).

2-(5-fluoro-1-(4-fluorobenzyl)-2-oxoindolin-3-ylidene)-*N*-*p*-tolylhydrazine carbothioamide (**5b**)

Yield 77%. mp 230–235°C. FT-IR ν_{max} cm^{-1} (KBr): 1126, 1287, 1681, 3054, 3143, 3212, 3251. ^1H -NMR δ : 2.35 (s,

3H, CH₃), 5.0 (s, 2H, CH₂), 7.0–7.8 (m, 11H, Ar–H), 10.87 (s, 1H, N–NH), 12.58 (s, 1H, NH).

2-(5-chloro-1-(4-methylbenzyl)-2-oxoindolin-3-ylidene)-N-(4-chlorophenyl)hydrazine carbothioamide (5c)

Yield 86%. mp 235–239°C. FT-IR ν_{\max} cm^{−1} (KBr): 1135, 1294, 1682, 3026, 3119, 3243, 3264. ¹H-NMR δ : 2.51 (s, 3H, CH₃), 5.01 (s, 2H, CH₂), 7.0–8.7 (m, 11H, Ar–H), 11.00 (s, 1H, N–NH), 12.58 (s, 1H, NH).

2-(5-chloro-1-(4-chlorobenzyl)-2-oxoindolin-3-ylidene)-N-(4-chlorophenyl)hydrazine carbothioamide (5d)

Yield 66%. mp 238–240°C. FT-IR ν_{\max} cm^{−1} (KBr): 1126, 1310, 1706, 3047, 3127, 3237, 3276. ¹H-NMR δ : 4.96 (s, 2H, CH₂), 7.0–7.9 (m, 11H, Ar–H), 10.99 (s, 1H, N–NH), 12.61 (s, 1H, NH).

2-(5-chloro-1-(4-fluorobenzyl)-2-oxoindolin-3-ylidene)-N-(4-chlorophenyl)hydrazine carbothioamide (5e)

Yield 81%. mp 230–235°C. FT-IR ν_{\max} cm^{−1} (KBr): 1156, 1295, 1696, 3056, 3123, 3219, 3259. ¹H-NMR δ : 5.00 (s, 2H, CH₂), 7.0–7.9 (m, 11H, Ar–H), 10.99 (s, 1H, N–NH), 12.59 (s, 1H, NH).

2-(5-bromo-1-(4-fluorobenzyl)-2-oxoindolin-3-ylidene)-N-(4-chlorophenyl)hydrazine carbothioamide (5f)

Yield 73%. mp 240–245°C. FT-IR ν_{\max} cm^{−1} (KBr): 1134, 1287, 1685, 3014, 3161, 3206, 3245. ¹H-NMR δ : 5.0 (s, 2H, CH₂), 7.0–8.0 (m, 11H, Ar–H), 11.0 (s, 1H, N–NH), 12.59 (s, 1H, NH).

2-(5-bromo-1-(4-fluorobenzyl)-2-oxoindolin-3-ylidene)-N-p-tolylhydrazine carbothioamide (5g)

Yield 80%. mp 268–270°C. FT-IR ν_{\max} cm^{−1} (KBr): 1125, 1309, 1694, 3081, 3123, 3204, 3261. ¹H-NMR δ : 2.37 (s, 3H, CH₃), 4.95 (s, 2H, CH₂), 6.9–8.0 (m, 11H, Ar–H), 10.92 (s, 1H, N–NH), 12.52 (s, 1H, NH).

2-(5-bromo-1-(4-methylbenzyl)-2-oxoindolin-3-ylidene)-N-(4-chlorophenyl)hydrazine carbothioamide (5h)

Yield 81%. mp 220–223°C. FT-IR ν_{\max} cm^{−1} (KBr): 1124, 1295, 1710, 3004, 3153, 3210, 3251. ¹H-NMR δ : 2.27 (s, 3H, CH₃), 4.96 (s, 2H, CH₂), 7.0–8.0 (m, 11H, Ar–H), 11.00 (s, 1H, N–NH), 12.61 (s, 1H, NH).

2-(5-bromo-1-(4-chlorobenzyl)-2-oxoindolin-3-ylidene)-N-p-tolylhydrazine carbothioamide (5i)

Yield 72%. mp 217–220°C. FT-IR ν_{\max} cm^{−1} (KBr): 1094, 1269, 1686, 3043, 3113, 3206, 3243. ¹H-NMR δ : 2.51 (s, 3H, CH₃), 3.35 (s, 2H, CH₂), 7.0–8.0 (m, 11H, Ar–H), 10.92 (s, 1H, N–NH), 12.51 (s, 1H, NH).

2-(5-bromo-1-(4-methylbenzyl)-2-oxoindolin-3-ylidene)-N-p-tolylhydrazine carbothioamide (5j)

Yield 79%. mp 185–190°C. FT-IR ν_{\max} cm^{−1} (KBr): 1126, 1309, 1698, 3056, 3133, 3218, 3239. ¹H-NMR δ : 2.27 (s, 3H, CH₃), 2.51 (s, 3H, CH₃), 4.96 (s, 2H, CH₂), 6.9–8.0 (m, 11H, Ar–H), 10.92 (s, 1H, N–NH), 12.54 (s, 1H, NH).

2-(5-bromo-1-(4-chlorobenzyl)-2-oxoindolin-3-ylidene)-N-(4-chlorophenyl)hydrazine carbothioamide (5k)

Yield 76%. mp 230–233°C. FT-IR ν_{\max} cm^{−1} (KBr): 1120, 1296, 1684, 3046, 3135, 3216, 3247. ¹H-NMR δ : 4.94 (s, 2H, CH₂), 6.4–7.7 (m, 11H, Ar–H), 11.29 (s, 1H, N–NH), 12.52 (s, 1H, NH).


Cytostatic assays

The methodology for undertaking the antiproliferative assays has been published previously (Baraldi *et al.*, 2004). In brief, varying concentrations of the compounds (5-fold dilutions) were incubated at 37°C for 72 h (Molt4/C8 and CEM T-lymphocytes) or 48 h (L1210 cells) in 200 μ l 96-well microtiter plates, and the viable tumor cell number was counted at the end of the incubation period using a Coulter Counter (Coulter Electronics, Harpenden Hertz, U.K.).

Results and discussion

Chemistry

The compounds in series **4** and **5** were prepared by the methodologies outlined in Scheme 1. The synthesis of *N*-benzyl-2,3-dioxindole (**3**) was carried out by reacting benzyl chloride with various 2,3-dioxo-indoles in the presence of potassium carbonate under reflux in DMF. The synthesis of the 2,3-dioxo-2,3-dihydroindolo-thiosemi-carbazones (**4a–g**, **5a–k**) was carried out by the condensation of various 2,3-dioxoindoles/*N*-arylalkyl-2,3-dioxoindoles (2,3) with aryl thiosemicarbazide under reflux in ethanol in the presence of catalytic amounts of glacial acetic acid. ¹H NMR spectroscopy indicated that the compounds exist as



Scheme 1 Synthesis of compounds **4a–g** and **5a–k**. The reagents used are as follows: *i* $\text{CCl}_3\text{CH}(\text{OH})_2/\text{H}_2\text{SO}_4/\text{Na}_2\text{SO}_4$; *ii* $\text{R}_3\text{-C}_6\text{H}_4\text{CH}_2\text{Cl}/\text{K}_2\text{CO}_3/\text{DMF}$; *iii* $\text{R}_4\text{-C}_6\text{H}_4\text{NHCSNHNH}_2$. The nature of the R_1 , R_2 , R_3 , and R_4 substituents are presented in Table 1

single isomers in solution. The compounds adopted the *Z* conformation (Karki *et al.*, 2009).

Biological activity

The compounds were evaluated for their cytostatic activity against human Molt4/C8 and CEM T-lymphocytes as well

as murine L1210 leukemia cells. The data are summarized in Table 1.

Several compounds (**4c–d** and **5b–c**) showed IC_{50} values in the low micromolar range (1–10 μM). Molt4/C8 cells were slightly more sensitive to the cytostatic activity of the compounds in both series **4** and **5** than CEM or L1210 cells.

Table 1 Cytostatic activity of test compounds against murine L1210 leukemia and human Molt4/C8 and CEM T-lymphocyte cells

Comp. code	Substituents				IC ₅₀ * (μM)		
	R ₁	R ₂	R ₃	R ₄	L1210	Molt4/C8	CEM
4a	Cl	H	H	Cl	108 ± 18	64 ± 20	61 ± 6
4b	Br	H	H	Cl	88 ± 39	29 ± 16	61 ± 11
4c	Br	H	H	CH ₃	9.6 ± 1.8	4.9 ± 3.5	6.8 ± 2.4
4d	F	H	H	CH ₃	8.2 ± 3.6	3.0 ± 0.7	4.8 ± 0.5
4e	F	Cl	H	Cl	69 ± 50	43 ± 10	48 ± 18
4f	F	Cl	H	CH ₃	166 ± 48	112 ± 81	132 ± 75
4g	F	H	H	Cl	34 ± 2	14 ± 7	25 ± 6
5a	F	H	F	Cl	154 ± 73	47 ± 3	56 ± 14
5b	F	H	F	CH ₃	6.2 ± 2.1	1.5 ± 1.1	2.5 ± 1.2
5c	Cl	H	CH ₃	Cl	9.8 ± 2.1	5.8 ± 4.2	18 ± 9
5d	Cl	H	Cl	Cl	124 ± 39	58 ± 13	68 ± 1
5e	Cl	H	F	Cl	103 ± 53	48 ± 5	50 ± 14
5f	Br	H	F	Cl	196 ± 66	48 ± 7	61 ± 9
5g	Br	H	F	CH ₃	190 ± 40	57 ± 9	132 ± 35
5h	Br	H	CH ₃	Cl	193 ± 49	74 ± 19	98 ± 14
5i	Br	H	Cl	CH ₃	≥ 500	288 ± 12	≥ 500
5j	Br	H	CH ₃	CH ₃	>500	393 ± 27	≥ 500
5k	Br	H	Cl	Cl	≥ 500	280 ± 43	392 ± 104
Melphalan	–	–	–	–	2.1 ± 0.02	3.2 ± 0.6	2.5 ± 0.2

* The IC₅₀ values represent the compound concentrations required to inhibit the growth of the tumor cells by 50%

There was, in general, a strong correlation between the three tumor cell lines regarding the cytostatic activities of the compounds. The most potent inhibitors of murine L1210 cell proliferation (i.e., **4c** and **4d**, and **5b** and **5c**) were also most inhibitory to human T-lymphocyte CEM and Molt4/C8 cell proliferation. Also, the scaffold of the most potent cytostatic compound in the **4** series (**4d**) was most inhibitory in the **5** series (**5b**). In the majority of cases, introduction of the R₃-benzyl substituent in the thiosemicarbazone derivative series **5** often led to (slightly) less potent inhibitors of tumor cell proliferation. In a number of cases, replacement of the chloro at R₄ by a methyl resulted in a marked potentiation of the cytostatic activity (compare **4b** with **4c**; **4g** with **4d**; **5a** with **5b**) (Table 1). In the **5** series of compounds, replacement of the chlorine atom at R₁ by a bromine often resulted in a decreased activity; i.e., compare **5c** with **5h**; **5d** with **5k**; and **5e** with **5f**.

Conclusion

Two series of derivatives of 2,3-dioxo-2,3-dihydroindol-(N-4-substituted aryl)-thiosemicarbazones were prepared in good yields. All compounds were evaluated for cytostatic

activity against human Molt4/C8 and CEM T-lymphocytes and murine L1210 leukemia cells. The compounds **4c**, **4d**, **5b**, and **5c** had IC₅₀ values in the low micromolar range (1–10 μM). Among the tested compounds, **5b** and **4d** had antiproliferative potencies that were close to those obtained for melphalan and should be the basis for further synthesis of more potent cytostatic agents.

Acknowledgments The study was supported by a grant from the K.U.Leuven (GOA no. 10/014). The authors want to thank Mrs. Lizzette van Berckelaer for excellent technical assistance.

References

- Azizian J, Shaidaei FB, Kefayati H (2003) A facile one pot method for the preparation of *N*-alkyl isatins under microwave irradiation. *Synth Commun* 33:789–793
- Baraldi PB, MDeLC Nunez, Tabrizo MA, De Clercq E, Balzarini J, Bermejo J, Estevez F, Romagnoli R (2004) Design, syntheses and biological evaluation of hybrid molecules containing α-methylene-γ-butyrolactones and polypyrrole minor groove binders. *J Med Chem* 47:2877–2886
- Karki SS, Mazumder UK, Gupta M, Bhattacharya S, Rathinasamy S, Thangavel S (2004) Synthesis, anticancer and antibacterial activity of some novel mononuclear Ru(II) complexes. *Chem Pharm Bull* 52:178–185
- Karki SS, Shrikanth T, Satyanarayana YD, Balzarini J, De Clercq E (2007) Synthesis, anticancer and cytotoxic activities of some mononuclear Ru(II) compounds. *Bioorg Med Chem* 15:6632–6641
- Karki SS, Bahaduria VS, Rana V, Kumar S, Prasanna GS, Das U, Balzarini J, De Clercq E, Dimmock JR (2009) 1-Arylmethyl-2,3-dioxo-2,3-dihydroindole thiosemicarbazones as leads for developing cytotoxins and anticonvulsants. *J Enzyme Inhib Med Chem* 24:537–544
- Lieber E, Pillai ECN, Hites RD (1957) Reactions of nitrous acid with 4-substituted thiosemicarbazides. *Can J Chem* 35:832–842
- Marvel CS, Heirs GS (1941) In Isatin. In: Blatt AH (ed) *Organic syntheses collection*, vol 1. Wiley, New York, pp 327–330
- Pandeya SN, Yogeeswari P, Sriram D, Nath G (1998) Synthesis and antimicrobial activity of *N*-Mannich bases of 3-[*N'*-sulphadioxo-imino]isatin and its methyl derivatives. *Bull Chim Farm* 137:321–324
- Pandeya SN, Yogeeswari P, Sriram D, De Clercq E, Pannecouque C, Witvrouw M (1999a) Synthesis and screening for anti-HIV activity of some *N*-Mannich bases of isatin derivatives. *Chemotherapy* 45:192–196
- Pandeya SN, Sriram D, Nath G, De Clercq E (1999b) Synthesis, antibacterial, antifungal and anti-HIV evaluation of Schiff and Mannich bases of isatin derivatives with 3-amino-2-methylmercapto quinazolin-4(3H)-one. *Pharm Acta Helv* 74:11–17
- Pandeya SN, Sriram D, Nath G, De Clercq E (2000) Synthesis, antibacterial, antifungal and anti-HIV activities of Norfloxacin Mannich bases. *Eur J Med Chem* 35:249–255
- Sarangapani M, Reddy VM (1994) Synthesis and antimicrobial activity of 1-[(*N,N*-disubstituted amino)methyl]-3-[(2-phenyl-3,4-dihydro-4-oxoquinazoline-3-yl)]indole-2-one. *Indian J Heterocycl Chem* 3:257–260
- Sen AB, Sengupta SK (1962) Possible antiamoebic agents. Part XIX. Synthesis of 1,3,4-thiadiazolines and di-1,3,4-thiadiazolines. *J Ind Chem Soc* 39:628–634
- Silva DJFM, Garden SJ, Pinto AC (2001) The chemistry of isatins: a review from 1975 to 1999. *J Braz Chem Soc* 12:273–324

- Singh SP, Shukla SK, Awasthi LP (1983) Synthesis of some 3-(4'-nitrobenzoylhydrazono)-2-indolinones as a potential antiviral agents. *Curr Sci* 52:766–769
- Somogyi L (2001) Transformation of isatin 3-acylhydrazones under acetylating conditions: synthesis and structure elucidation of 1,5'-disubstituted 3'-acetylspiro[oxindole-3,2'-[1,3,4]oxadiazolines]. *Bull Chem Soc Jpn* 74:873–881
- Sridhar SK, Pandeya SN, Stables JP, Ramesh A (2002) The wide pharmacological versatility of semicarbazones, thiosemicarbazones and their metal complexes. *Eur J Med Chem* 16:129–132
- Sriram D, Pandeya SN, Nath G, De Clercq E (2000) Synthesis, antibacterial, antifungal and anti-HIV evaluation of Schiff and Mannich bases of isatin and its derivatives with triazole. *Arzneimittelforschung* 50:55–59
- Varma RS, Nobles WL (1975) Antiviral, antibacterial, and antifungal activities of isatin *N*-Mannich bases. *J Pharm Sci* 64:881–882
- Varma M, Pandeya SN, Singh KN, Stables JP (2004) Anticonvulsant activity of Schiff bases of isatin derivatives. *Acta Pharm* 54:49–56